

## **Tachinidae Parasitoids of *Traumatocampa ispartaensis* from Turkey**

Mustafa Avcı<sup>1</sup> and Kenan Kara<sup>2</sup>

Tachinid parasitoids of *Traumatocampa ispartaensis* Doğanlar & Avcı (Lepidoptera, Thaumetopoeidae), which was found to be a new species, were collected from the cedar forests around Isparta-Kapıdağ. The species found were *Blondelia nigripes* (Fall.), *Compsilura concinnata* (Meig.), *Pales processioneae* (Ratz.), *Phryxe caudata* (Rond.), *Exorista segregata* (Rond.) and *Carcelia iliaca* (Ratz.). Within the six species of Tachinidae, *B. nigripes* was the most common one, parasitizing up to 4.6% of *T. ispartaensis* pupae.

**KEY WORDS:** Parasitoid; Tachinidae; *Traumatocampa ispartaensis*; *Blondelia nigripes*; Turkey.

---

Received Sept. 7, 2001; received in final form March 11, 2002; <http://www.phytoparasitica.org> posting July 2, 2002.

<sup>1</sup>Süleyman Demirel University, Faculty of Forestry, Forest Entomology and Protection Dept., Isparta, Turkey [Fax: +90-246-2371810; e-mail: [avci@orman.sdu.edu.tr](mailto:avci@orman.sdu.edu.tr)].

<sup>2</sup>Gaziosmanpaşa University, Faculty of Agriculture, Plant Protection Dept., Tokat, Turkey [Fax: +90-356-2521488; e-mail: [kkara@mail.gop.edu.tr](mailto:kkara@mail.gop.edu.tr)].