

Tachinid Parasitoids (Diptera: Tachinidae) of *Spodoptera exigua* in Cotton Fields in Diyarbakır, Turkey

Levent Efil*,¹ and Kenan Kara²

Tachinid parasitoids *Exorista xanthaspis* (Wiedemann), *Nemorilla maculosa* (Meigen), *Palesisa maculosa* (Villeneuve) and *Drino imberbis* (Wiedemann) were obtained from *Spodoptera exigua* (Hübner) (Noctuidae) larvae collected from cotton fields in Turkey. *S. exigua* is a new host record for *N. maculosa*, and *P. maculosa* is recorded from Turkey for the first time.

KEY WORDS: Parasitoids; Tachinidae; *Spodoptera exigua* (Hübner); *Exorista xanthaspis* (Wiedemann); *Nemorilla maculosa* (Meigen); *Palesisa maculosa* (Villeneuve); *Drino imberbis* (Wiedemann); Turkey.

Received Oct. 16, 2003; accepted Dec. 23, 2003; <http://www.phytoparasitica.org> posting July 14, 2004.

¹Plant Protection Research Institute, Diyarbakır, Turkey. *Corresponding author [Fax: +90-412-2234775; e-mail: efil46@hotmail.com].

²Dept. of Plant Protection, Faculty of Agriculture, Gaziosmanpaşa University, Tokat, Turkey [e-mail: kkara@gop.edu.tr].